

Chapter 1

Recursion

File: `set-theory/sections/04_recursion.ftl.tex`

[readtex `set-theory/sections/02_ordinals.ftl.tex`]

SET_THEORY_04_7107446027845632

Definition 1.1. Let A be a class and α be an ordinal.

$$A^{<\alpha} = \{f \mid f \text{ is a map from } \beta \text{ to } A \text{ for some ordinal } \beta \text{ less than } \alpha\}.$$

SET_THEORY_04_1955917673267200

Definition 1.2. Let A be a class.

$$A^{<\infty} = \{f \mid f \text{ is a map from } \alpha \text{ to } A \text{ for some ordinal } \alpha\}.$$

SET_THEORY_04_7841726894964736

Lemma 1.3. Let A be a class and f be a map to A such that $\text{dom}(f)$ is a transitive subclass of **Ord** and $\alpha \in \text{dom}(f)$. Then $f \upharpoonright \alpha \in A^{<\infty}$.

SET_THEORY_04_5597213870784512

Definition 1.4. Let H be a map and $G : A^{<\infty} \rightarrow A$ for some class A such that H is a map to A . H is recursive regarding G iff $\text{dom}(H)$ is a transitive subclass of **Ord** and for all $\alpha \in \text{dom}(H)$ we have

$$H(\alpha) = G(H \upharpoonright \alpha).$$

SET_THEORY_04_2876133366300672

Proposition 1.5. Let A be a class and G be a map from $A^{<\infty}$ to A . Let H, H' be maps to A that are recursive regarding G . Then

$$H(\alpha) = H'(\alpha)$$

for all $\alpha \in \text{dom}(H) \cap \text{dom}(H')$.

Proof. Define $\Phi = \{\alpha \in \mathbf{Ord} \mid \text{if } \alpha \in \text{dom}(H) \cap \text{dom}(H') \text{ then } H(\alpha) = H'(\alpha)\}$.

For all ordinals α if every ordinal less than α lies in Φ then $\alpha \in \Phi$.

Proof. Let $\alpha \in \mathbf{Ord}$. Assume that every $y \in \alpha$ lies in Φ .

Let us show that if $\alpha \in \text{dom}(H) \cap \text{dom}(H')$ then $H(\alpha) = H'(\alpha)$. Suppose $\alpha \in \text{dom}(H) \cap \text{dom}(H')$. Then $\alpha \subseteq \text{dom}(H), \text{dom}(H')$. Indeed $\text{dom}(H)$ and $\text{dom}(H')$ are transitive classes. Hence for all $y \in \alpha$ we have $y \in \text{dom}(H) \cap \text{dom}(H')$. Thus $H(y) = H'(y)$ for all $y \in \alpha$. Therefore $H \upharpoonright \alpha = H' \upharpoonright \alpha$. H and H' are recursive regarding G . Hence $H(\alpha) = G(H \upharpoonright \alpha) = G(H' \upharpoonright \alpha) = H'(\alpha)$. End.

Thus $\alpha \in \Phi$. Qed.

[prover vampire] Then Φ contains every ordinal (by ??). Therefore we have $H(\alpha) = H'(\alpha)$ for all $\alpha \in \text{dom}(H) \cap \text{dom}(H')$. \square

SET_THEORY_04_3600210873810944

Theorem 1.6 (Recursion theorem). Let A be a class and G be a map from $A^{<\infty}$ to A . Then there exists a map F from **Ord** to A that is recursive regarding G .

Proof. Every ordinal is contained in the domain of some map H to A such that H is recursive regarding G .

Proof. Define

$$\Phi = \left\{ \alpha \in \mathbf{Ord} \mid \alpha \text{ is contained in the domain of some map to } A \text{ that is recursive regarding } G \right\}.$$

Let us show that for every ordinal α if every ordinal less than α lies in Φ then $\alpha \in \Phi$. Let α be an ordinal. Assume that every ordinal less than α lies in Φ . Then for all $y \in \alpha$ there exists a map h to A such that h is recursive regarding G and $y \in \text{dom}(h)$. Define $H'(y) =$ “choose a map h to A such that h is recursive regarding G and $y \in \text{dom}(h)$ in $h(y)$ ” for $y \in \alpha$. Then H' is a map from α to A . We have $H' = H' \upharpoonright \alpha$. Define

$$H(\beta) = \begin{cases} H'(\beta) & : \beta < \alpha \\ G(H' \upharpoonright \beta) & : \beta = \alpha \end{cases}$$

for $\beta \in \text{succ}(\alpha)$. Then $H \upharpoonright \beta \in A^{<\infty}$ for all $\beta \in \text{dom}(H)$.

(a) $\text{dom}(H)$ is a transitive subclass of **Ord**.

(b) For all $\beta \in \text{dom}(H)$ we have $H(\beta) = G(H \upharpoonright \beta)$.

Proof. Let $\beta \in \text{dom}(H)$. Then $\beta < \alpha$ or $\beta = \alpha$.

Case $\beta < \alpha$. Choose a map h to A such that h is recursive regarding G and $\beta \in \text{dom}(h)$ and $H'(\beta) = h(\beta)$.

Let us show that for all $y \in \beta$ we have $h(y) = H(y)$. Let $y \in \beta$. Then $H(y) = H'(y)$. Choose a map h' to A such that h' is recursive regarding G and $y \in \text{dom}(h')$ and $H'(y) = h'(y)$. [prover vampire] Then $h'(y) = h(y)$ (by proposition 1.5). Indeed $y \in \text{dom}(h) \cap \text{dom}(h')$. End.

Hence $h \upharpoonright \beta = H \upharpoonright \beta$. Thus $H(\beta) = H'(\beta) = h(\beta) = G(h \upharpoonright \beta) = G(H \upharpoonright \beta)$. End.

Case $\beta = \alpha$. We have $H \upharpoonright \alpha = H' \upharpoonright \alpha$. End. Qed.

Hence H is a map to A such that H is recursive regarding G and $\alpha \in \text{dom}(H)$. Thus $\alpha \in \Phi$. End.

[prover vampire] Therefore Φ contains every ordinal (by ??). Consequently every ordinal is contained in the domain of some map H to A such that H is recursive regarding G . Qed.

Define $F(\alpha) =$ “choose a map H to A such that H is recursive regarding G and $\alpha \in \text{dom}(H)$ in $H(\alpha)$ ” for $\alpha \in \mathbf{Ord}$. Then F is a map from **Ord** to A .

F is recursive regarding G .

Proof. (a) $\text{dom}(F)$ is a transitive subclass of **Ord**.

(b) For all $\alpha \in \mathbf{Ord}$ we have $F(\alpha) = G(F \upharpoonright \alpha)$.

Proof. Let $\alpha \in \mathbf{Ord}$. Choose a map H to A such that H is recursive regarding G and $\alpha \in \text{dom}(H)$ and $F(\alpha) = H(\alpha)$.

Let us show that $F(\beta) = H(\beta)$ for all $\beta \in \alpha$. Let $\beta \in \alpha$. Choose a map H' to A such that H' is recursive regarding G and $\beta \in \text{dom}(H')$ and $F(\beta) = H'(\beta)$. [prover vampire] Then $H(\beta) = H'(\beta)$ (by proposition 1.5). Indeed $\beta \in \text{dom}(H) \cap \text{dom}(H')$. Therefore $F(\beta) = H'(\beta)$. End.

Hence $H \upharpoonright \alpha = F \upharpoonright \alpha$. Thus $F(\alpha) = H(\alpha) = G(H \upharpoonright \alpha) = G(F \upharpoonright \alpha)$. Qed. Qed. \square

SET_THEORY_04_1787371569807360

Theorem 1.7. Let A be a class and G be a map from $A^{<\infty}$ to A . Let F, F' be maps from **Ord** to A that are recursive regarding G . Then $F = F'$.

Proof. F and F' are recursive regarding G . [prover vampire] Then $F(\alpha) = F'(\alpha)$ for all $\alpha \in \text{dom}(F) \cap \text{dom}(F')$ (by proposition 1.5). Indeed let $\alpha \in \text{dom}(F) \cap \text{dom}(F')$. We have $\text{dom}(F) = \mathbf{Ord} = \text{dom}(F')$. Hence $F(\alpha) = F'(\alpha)$ for all $\alpha \in \mathbf{Ord}$. Thus $F = F'$. \square

SET_THEORY_04_8446954438656000

Theorem 1.8. Let A be a class. Let $a \in A$ and $G : \mathbf{Ord} \times A \rightarrow A$ and $H : \mathbf{Ord} \times A^{<\infty} \rightarrow A$. Then there exists a map F from **Ord** to A such that

$$F(0) = a$$

and for all ordinals α we have

$$F(\text{succ}(\alpha)) = G(\alpha, F(\alpha))$$

and for all limit ordinals λ we have

$$F(\lambda) = H(\lambda, F \upharpoonright \lambda).$$

Proof. Define

$$J(f) = \begin{cases} a & : \text{dom}(f) = 0 \\ G(\text{pred}(\text{dom}(f)), f(\text{pred}(\text{dom}(f)))) & : \text{dom}(f) \text{ is a successor ordinal} \\ H(\text{dom}(f), f) & : \text{dom}(f) \text{ is a limit ordinal} \end{cases}$$

for $f \in A^{<\infty}$.

Then J is a map from $A^{<\infty}$ to A . Indeed we can show that for any $f \in A^{<\infty}$ we have $J(f) \in A$. Let $f \in A^{<\infty}$. Take $\alpha \in \mathbf{Ord}$ such that $f : \alpha \rightarrow A$. If $\alpha = 0$ then $J(f) = a \in A$. If α is a successor ordinal then $J(f) = G(\text{pred}(\alpha), f(\text{pred}(\alpha))) \in A$. If α is a limit ordinal then $J(f) = H(\alpha, f) \in A$. End.

Hence we can take a map F from **Ord** to A that is recursive regarding J . Then $F \upharpoonright \alpha \in A^{<\infty}$ for any ordinal α .

(1) $F(0) = a$.

Proof. $F(0) = J(F \upharpoonright 0) = a$. Qed.

(2) $F(\text{succ}(\alpha)) = G(\alpha, F(\alpha))$ for all ordinals α .

Proof. Let α be an ordinal. Then $F(\text{succ}(\alpha)) = J(F \upharpoonright \text{succ}(\alpha)) = G(\text{pred}(\text{succ}(\alpha)), (F \upharpoonright \text{succ}(\alpha))(\text{pred}(\text{succ}(\alpha)))) = G(\alpha, (F \upharpoonright \text{succ}(\alpha))(\alpha)) = G(\alpha, F(\alpha))$. Qed.

(3) $F(\lambda) = H(\lambda, F \upharpoonright \lambda)$ for all limit ordinals λ .

Proof. Let λ be a limit ordinal. Then $F(\lambda) = J(F \upharpoonright \lambda) = H(\lambda, F \upharpoonright \lambda)$. Qed. \square